

Effects of exercise training on patients with CKD 2018

- 1: Afsar B, Siriopol D, Aslan G, Eren OC, Dagele T, Kilic U, Kanbay A, Burlacu A, Covic A, Kanbay M. The impact of exercise on physical function, cardiovascular outcomes and quality of life in chronic kidney disease patients: a systematic review. *Int Urol Nephrol*. 2018 May;50(5):885-904. doi: 10.1007/s11255-018-1790-4. Epub 2018 Jan 17. Review.
- 2: Aoike DT, Baria F, Kamimura MA, Ammirati A, Cuppari L. Home-based versus center-based aerobic exercise on cardiopulmonary performance, physical function, quality of life and quality of sleep of overweight patients with chronic kidney disease. *Clin Exp Nephrol*. 2018 Feb;22(1):87-98. doi: 10.1007/s10157-017-1429-2.
- 3: Barcellos FC, Del Vecchio FB, Reges A, Mielke G, Santos IS, Umpierre D, Bohlke M, Hallal PC. Exercise in patients with hypertension and chronic kidney disease: a randomized controlled trial. *J Hum Hypertens*. 2018 Jun;32(6):397-407. doi: 10.1038/s41371-018-0055-0.
- 4: Beetham KS, Howden EJ, Krishnasamy R, Isbel NM, Coombes JS. Feasibility of higher intensity exercise in patients with chronic kidney disease. *J Sports Med Phys Fitness*. 2018 Jan-Feb;58(1-2):127-134. doi: 10.23736/S0022-4707.16.06575-0.
- 5: Beetham KS, Howden EJ, Isbel NM, Coombes JS. Agreement between cystatin-C and creatinine based eGFR estimates after a 12-month exercise intervention in patients with chronic kidney disease. *BMC Nephrol*. 2018 Dec 18;19(1):366. doi: 10.1186/s12882-018-1146-4.
- 6: Bohm CJ, Storsley LJ, Hiebert BM, Nelko S, Tangri N, Cheskin LJ, McAdams-DeMarco MA, Rigatto C. Impact of Exercise Counseling on Physical Function in Chronic Kidney Disease: An Observational Study. *Can J Kidney Health Dis*. 2018 Feb 19;5:2054358117753615. doi: 10.1177/2054358117753615.
- 7: Fuzari HK, Dornelas de Andrade A, A Rodrigues M, I Medeiros A, F Pessoa M, Lima AM, Cerqueira MS, Marinho PE. Whole body vibration improves maximum voluntary isometric contraction of knee extensors in patients with chronic kidney disease: A randomized controlled trial. *Physiother Theory Pract*. 2018 Feb 26:1-10. doi: 10.1080/09593985.2018.1443537.

8: Greenwood SA, Castle E, Lindup H, Mayes J, Waite I, Grant D, Mangahis E, Crabb O, Shevket K, Macdougall IC, MacLaughlin HL. Mortality and morbidity following exercise-based renal rehabilitation in patients with chronic kidney disease: the effect of programme completion and change in exercise capacity. *Nephrol Dial Transplant*. 2018 Nov 30. doi: 10.1093/ndt/gfy351.

9: Hellberg M, Höglund P, Svensson P, Clyne N. Comparing effects of 4 months of two self-administered exercise training programs on physical performance in patients with chronic kidney disease: RENEXC - A randomized controlled trial. *PLoS One*. 2018 Dec 20;13(12):e0207349. doi: 10.1371/journal.pone.0207349.

10: Highton PJ, Neale J, Wilkinson TJ, Bishop NC, Smith AC. Physical activity, immune function and inflammation in kidney patients (the PINK study): a feasibility trial protocol. *BMJ Open*. 2017 May 29;7(5):e014713. doi: 10.1136/bmjopen-2016-014713.

11: Ikizler TA, Robinson-Cohen C, Ellis C, Headley SAE, Tuttle K, Wood RJ, Evans EE, Milch CM, Moody KA, Germain M, Limkunakul C, Bian A, Stewart TG, Himmelfarb J. Metabolic Effects of Diet and Exercise in Patients with Moderate to Severe CKD: A Randomized Clinical Trial. *J Am Soc Nephrol*. 2018 Jan;29(1):250-259. doi: 10.1681/ASN.2017010020.

12: Jagannathan R, Ziolkowski SL, Weber MB, Cobb J, Pham N, Long J, Anand S, Lobelo F. Physical activity promotion for patients transitioning to dialysis using the "Exercise is Medicine" framework: a multi-center randomized pragmatic trial (EIM-CKD trial) protocol. *BMC Nephrol*. 2018 Sep 12;19(1):230. doi: 10.1186/s12882-018-1032-0.

13: MacKinnon HJ, Wilkinson TJ, Clarke AL, Gould DW, O'Sullivan TF, Xenophontos S, Watson EL, Singh SJ, Smith AC. The association of physical function and physical activity with all-cause mortality and adverse clinical outcomes in nondialysis chronic kidney disease: a systematic review. *Ther Adv Chronic Dis*. 2018 Jul 4;9(11):209-226. doi: 10.1177/2040622318785575.

- 14: Martini A, Ammirati A, Garcia C, Andrade C, Portela O, Cendoroglo MS, Sesso R. Evaluation of quality of life, physical, and mental aspects in longevous patients with chronic kidney disease. *Int Urol Nephrol*. 2018 Apr;50(4):725-731. doi: 10.1007/s11255-018-1813-1.
- 15: Nishiwaki H, Hasegawa T, Shinji M, Matsuo F, Watanabe T, Makino H, Koiwa F, Hishida A. Practice pattern of physician's directions of exercise restriction in patients with chronic kidney disease: results from the Chronic Kidney Disease Japan Cohort study. *Clin Exp Nephrol*. 2018 Oct;22(5):1108-1115. doi: 10.1007/s10157-018-1562-6.
- 16: Souza MK, Neves RVP, Rosa TS, Cenedeze MA, Arias SCA, Fujihara CK, Bacurau RFP, Câmara NOS, Moraes MR, Pacheco E Silva Filho A. Resistance training attenuates inflammation and the progression of renal fibrosis in chronic renal disease. *Life Sci*. 2018 Aug 1;206:93-97. doi: 10.1016/j.lfs.2018.05.034.
- 17: Vanden Wyngaert K, Van Craenenbroeck AH, Van Biesen W, Dhondt A, Tanghe A, Van Ginckel A, Celie B, Calders P. The effects of aerobic exercise on eGFR, blood pressure and VO₂peak in patients with chronic kidney disease stages 3-4: A systematic review and meta-analysis. *PLoS One*. 2018 Sep 11;13(9):e0203662. doi: 10.1371/journal.pone.0203662
- 18: Wallin H, Asp AM, Wallquist C, Jansson E, Caidahl K, Hylander Rössner B, Jacobson SH, Rickenlund A, Eriksson MJ. Gradual reduction in exercise capacity in chronic kidney disease is associated with systemic oxygen delivery factors. *PLoS One*. 2018 Dec 19;13(12):e0209325. doi: 10.1371/journal.pone.0209325
- 19: Watson EL, Gould DW, Wilkinson TJ, Xenophontos S, Clarke AL, Vogt BP, Viana JL, Smith AC. Twelve-week combined resistance and aerobic training confers greater benefits than aerobic training alone in nondialysis CKD. *Am J Physiol Renal Physiol*. 2018 Jun 1;314(6):F1188-F1196. doi: 10.1152/ajprenal.00012.2018.
- 20: Wilkinson TJ, Watson EL, Xenophontos S, Gould DW, Smith AC. The 'minimum clinically important difference' in frequently reported objective physical function tests following a 12-week renal rehabilitation exercise intervention in non-dialysis chronic kidney disease. *Am J Phys Med Rehabil*. 2018 Oct 22. doi: 10.1097/PHM.0000000000001080.

21: Wilkinson TJ, Gould DW, Nixon DGD, Watson EL, Smith AC. Quality over quantity? Association of skeletal muscle myosteatosis and myofibrosis on physical function in chronic kidney disease. *Nephrol Dial Transplant*. 2018 Jun 22. doi: 10.1093/ndt/gfy139.

22: Wilkinson TJ PhD, Xenophontos S MSc, Gould DW PhD, Vogt BP PhD, Viana JL PhD, Smith AC PhD, Watson EL PhD. Test-retest reliability, validation, and "minimal detectable change" scores for frequently reported tests of objective physical function in patients with non-dialysis chronic kidney disease. *Physiother Theory Pract*. 2018 Mar 30;1-12. doi: 10.1080/09593985.2018.1455249

Effects of exercise training on dialysis patients 2018

1: Aragoncillo I, Ligerio JM, Hevia C, Morales AL, Amézquita Y, Cervera T, Vega A, Abad S, Macías N, Luño J. Rationale and design of the PHYSICALFAV trial: a randomized controlled trial to evaluate the effect of preoperative isometric exercise on vascular calibre and maturation of autologous arteriovenous fistulas. *Clin Kidney J*. 2018 Dec;11(6):841-845. doi: 10.1093/ckj/sfy046.

2: Baggetta R, D'Arrigo G, Torino C, ElHafeez SA, Manfredini F, Mallamaci F, Zoccali C, Tripepi G; EXCITE Working group. Effect of a home based, low intensity, physical exercise program in older adults dialysis patients: a secondary analysis of the EXCITE trial. *BMC Geriatr*. 2018 Oct 20;18(1):248. doi: 10.1186/s12877-018-0938-5.

3: Brown PDS, Rowed K, Shearer J, MacRae JM, Parker K. Impact of intradialytic exercise intensity on urea clearance in hemodialysis patients. *Appl Physiol Nutr Metab*. 2018 Jan;43(1):101-104. doi: 10.1139/apnm-2017-0460.

4: Bučar Pajek M, Pajek J. Characterization of deficits across the spectrum of motor abilities in dialysis patients and the impact of sarcopenic overweight and obesity. *Clin Nutr*. 2018 Jun;37(3):870-877. doi: 10.1016/j.clnu.2017.03.008.

5: Cho JH, Lee JY, Lee S, Park H, Choi SW, Kim JC. Effect of intradialytic exercise on daily physical activity and sleep quality in maintenance hemodialysis patients. *Int Urol Nephrol*. 2018 Apr;50(4):745-754. doi: 10.1007/s11255-018-1796-y.

6: Chou HY, Chen SC, Yen TH, Han HM. Effect of a Virtual Reality-Based Exercise Program on Fatigue in Hospitalized Taiwanese End-Stage Renal Disease Patients Undergoing Hemodialysis. *Clin Nurs Res*. 2018 Jul 1;1054773818788511. doi: 10.1177/1054773818788511

7: Cooke AB, Ta V, Iqbal S, Gomez YH, Mavrakanas T, Barré P, Vasilevsky M, Rahme E, Daskalopoulou SS. The Impact of Intradialytic Pedaling Exercise on Arterial Stiffness: A Pilot Randomized Controlled Trial in a Hemodialysis Population. *Am J Hypertens*. 2018 Mar 10;31(4):458-466. doi: 10.1093/ajh/hpx191.

8: Esgalhado MGBM, Stockler-Pinto MB, Cardozo LFMF, Barboza JE, Mafra D. Does high intensity exercise affects irisin plasma levels in hemodialysis patients? A pilot study. *J Bras Nefrol*. 2018 Jan-Mar;40(1):53-58. doi: 10.1590/1678-4685-JBN-3802.

9: Frih B, Mkacher W, Jaafar H, Frih A, Ben Salah Z, El May M, Hammami M. Specific balance training included in an endurance-resistance exercise program improves postural balance in elderly patients undergoing haemodialysis. *Disabil Rehabil*. 2018 Apr;40(7):784-790. doi: 10.1080/09638288.2016.1276971.

10: Fuhro MI, Dorneles GP, Andrade FP, Romão PRT, Peres A, Monteiro MB. Acute exercise during hemodialysis prevents the decrease in natural killer cells in patients with chronic kidney disease: a pilot study. *Int Urol Nephrol*. 2018 Mar;50(3):527-534. doi: 10.1007/s11255-017-1747-z.

11: García Testal A, García Maset R, Hervás Marín D, Pérez-Domínguez B, Royo Maicas P, Rico Salvador IS, Meléndez-Oliva E, Molina Aracil J, Murgui Chiva M, Del Pozo Blanco O, Olagüe Díaz P, Fernández Najera JE, Torregrosa De Juan E, Benedito Carrera C, Segura-Ortí E. Influence of Physical Exercise on the Dialytic Adequacy Parameters of Patients on Hemodialysis. *Ther Apher Dial*. 2018 Sep 18. doi: 10.1111/1744-9987.12762.

12: Gomes Neto M, de Lacerda FFR, Lopes AA, Martinez BP, Saquetto MB. Intradialytic exercise training modalities on physical functioning and health-related quality of life in patients undergoing maintenance hemodialysis: systematic review and meta-analysis. *Clin Rehabil*. 2018 Sep;32(9):1189-1202.

13: Han M, Ye X, Preciado P, Williams S, Campos I, Bonner M, Young C, Marsh D, Larkin JW, Usvyat LA, Maddux FW, Pecoits-Filho R, Kotanko P. Relationships between Neighborhood Walkability and Objectively Measured Physical Activity Levels in Hemodialysis Patients. *Blood Purif.* 2018;45(1-3):236-244. doi: 10.1159/000485161.

14: Hatef M, Sharif Nia H, Boyle C, Shafipour V. The Validity and Reliability of the Exercise Self-Efficacy Scale in a Sample of Hemodialysis Patients. *J Nurs Meas.* 2018 Dec;26(3):566-578. doi: 10.1891/1061-3749.26.3.566.

15: Jagannathan R, Ziolkowski SL, Weber MB, Cobb J, Pham N, Long J, Anand S, Lobelo F. Physical activity promotion for patients transitioning to dialysis using the "Exercise is Medicine" framework: a multi-center randomized pragmatic trial (EIM-CKD trial) protocol. *BMC Nephrol.* 2018 Sep 12;19(1):230. doi: 10.1186/s12882-018-1032-0.

16: Jayaseelan G, Bennett PN, Bradshaw W, Wang W, Rawson H. Exercise Benefits and Barriers: The Perceptions of People Receiving Hemodialysis. *Nephrol Nurs J.* 2018 Mar-Apr;45(2):185-219.

17: Jeong JH, Biruete A, Fernhall B, Wilund KR. Effects of acute intradialytic exercise on cardiovascular responses in hemodialysis patients. *Hemodial Int.* 2018 Oct;22(4):524-533. doi: 10.1111/hdi.12664

18: Kim JS, Yi JH, Shin J, Kim YS, Han SW. Effect of acute intradialytic aerobic and resistance exercise on one-day blood pressure in patients undergoing hemodialysis: a pilot study. *J Sports Med Phys Fitness.* 2018 Feb 26. doi: 10.23736/S0022-4707.18.07921-5

19: Kontos P, Grigorovich A, Colobong R, Miller KL, Nesrallah GE, Binns MA, Alibhai SMH, Parsons T, Jassal SV, Thomas A, Naglie G. Fit for Dialysis: a qualitative exploration of the impact of a research-based film for the promotion of exercise in hemodialysis. *BMC Nephrol.* 2018 Aug 6;19(1):195. doi: 10.1186/s12882-018-0984-4.

20: Martin N, Smith AC, Dungey MR, Young HML, Burton JO, Bishop NC. Exercise during hemodialysis does not affect the phenotype or prothrombotic nature of microparticles but alters their proinflammatory function. *Physiol Rep*. 2018 Sep;6(19):e13825. doi: 10.14814/phy2.13825.

21: Matsuzawa R, Roshanravan B, Shimoda T, Mamorita N, Yoneki K, Harada M, Watanabe T, Yoshida A, Takeuchi Y, Matsunaga A. Physical Activity Dose for Hemodialysis Patients: Where to Begin? Results from a Prospective Cohort Study. *J Ren Nutr*. 2018 Jan;28(1):45-53. doi: 10.1053/j.jrn.2017.07.004.

22: Matsuzawa R, Roshanravan B. Management of Physical Frailty in Patients Requiring Hemodialysis Therapy. *Contrib Nephrol*. 2018;196:101-109. doi: 10.1159/000485707.

23: McAdams-DeMarco MA, Konel J, Warsame F, Ying H, González Fernández M, Carlson MC, Fine DM, Appel LJ, Segev DL. Intradialytic Cognitive and Exercise Training May Preserve Cognitive Function. *Kidney Int Rep*. 2017 Aug 19;3(1):81-88. doi: 10.1016/j.ekir.2017.08.006.

24: McGregor G, Ennis S, Powell R, Hamborg T, Raymond NT, Owen W, Aldridge N, Evans G, Goodby J, Hewins S, Banerjee P, Krishnan NS, Ting SMS, Zehnder D. Feasibility and effects of intra-dialytic low-frequency electrical muscle stimulation and cycle training: A pilot randomized controlled trial. *PLoS One*. 2018 Jul 11;13(7):e0200354. doi: 10.1371/journal.pone.0200354

25: Medeiros AIC, Brandão DC, Souza RJP, Fuzari HKB, Barros CESR, Barbosa JBN, Leite JC, Cavalcanti FCB, Dornelas de Andrade A, de Melo Marinho PÉ. Effects of daily inspiratory muscle training on respiratory muscle strength and chest wall regional volumes in haemodialysis patients: a randomised clinical trial. *Disabil Rehabil*. 2018 Jul 27:1-8. doi: 10.1080/09638288.2018.1485181

26: Molina-Robles E, Colomer-Codinachs M, Roquet-Bohils M, Chirveches-Pérez E, Ortiz-Jurado P, Subirana-Casacuberta M. Effectiveness of an educational intervention and physical exercise on the functional capacity of patients on haemodialysis. *Enferm Clin*. 2018 May - Jun;28(3):162-170. doi: 10.1016/j.enfcli.2017.12.003.

27: Paluchamy T, Vaidyanathan R. Effectiveness of intradialytic exercise on dialysis adequacy, physiological parameters, biochemical markers and quality of life - A pilot study. *Saudi J Kidney Dis Transpl.* 2018 Jul-Aug;29(4):902-910. doi: 10.4103/1319-2442.239661.

28: Reboredo MM, Neder JA, Pinheiro BV, Henrique DM, Faria RS, Paula RB. Constant work-rate test to assess the effects of intradialytic aerobic training in mildly impaired patients with end-stage renal disease: a randomized controlled trial. *Arch Phys Med Rehabil.* 2011 Dec;92(12):2018-24. doi: 10.1016/j.apmr.2011.07.190.

29: Regolisti G, Maggiore U, Sabatino A, Gandolfini I, Pioli S, Torino C, Aucella F, Cupisti A, Pistolesi V, Capitanini A, Caloro G, Gregorini M, Battaglia Y, Mandreoli M, Dani L, Mosconi G, Bellizzi V, Di Iorio BR, Conti P, Fiaccadori E; Gruppo di Studio "Esercizio fisico nel paziente con insufficienza renale cronica" of the Società Italiana di Nefrologia. Interaction of healthcare staff's attitude with barriers to physical activity in hemodialysis patients: A quantitative assessment. *PLoS One.* 2018 Apr 27;13(4):e0196313. doi: 10.1371/journal.pone.0196313.

30: Rosa CSDC, Nishimoto DY, Souza GDE, Ramirez AP, Carletti CO, Daibem CGL, Sakkas GK, Monteiro HL. Effect of continuous progressive resistance training during hemodialysis on body composition, physical function and quality of life in end-stage renal disease patients: a randomized controlled trial. *Clin Rehabil.* 2018 Jul;32(7):899-908. doi: 10.1177/0269215518760696.

31: Salhab N, Karavetian M, Kooman J, Fiaccadori E, El Khoury CF. Effects of intradialytic aerobic exercise on hemodialysis patients: a systematic review and meta-analysis. *J Nephrol.* 2019 Jan 18. doi: 10.1007/s40620-018-00565-z

32: Shie JR, Chen TY, Kao CW. [The Effect of Exercise Training on Heart Rate Variability in Patients With Hemodialysis: A Systematic Review]. *Hu Li Za Zhi.* 2019 Feb;66(1):70-83. doi: 10.6224/JN.201902_66(1).09. Chinese

33: Shimizu U, Aoki H, Sakagami M, Akazawa K. Walking ability, anxiety and depression, significantly decrease EuroQol 5-Dimension 5-Level scores in older hemodialysis patients in Japan. *Arch Gerontol Geriatr.* 2018 Sep - Oct;78:96-100. doi: 10.1016/j.archger.2018.06.006.

34: Song YY, Hu RJ, Diao YS, Chen L, Jiang XL. Effects of Exercise Training on Restless Legs Syndrome, Depression, Sleep Quality, and Fatigue Among Hemodialysis Patients: A Systematic Review and Meta-analysis. *J Pain Symptom Manage*. 2018 Apr;55(4):1184-1195. doi: 10.1016/j.jpainsymman.2017.12.472.

35: Souweine JS, Kuster N, Chenine L, Rodriguez A, Patrier L, Morena M, Badia E, Chalabi L, Raynal N, Ohresser I, Leray-Moragues H, Mercier J, Hayot M, Le Quintrec M, Gouzi F, Cristol JP. Physical inactivity and protein energy wasting play independent roles in muscle weakness in maintenance haemodialysis patients. *PLoS One*. 2018 Aug 1;13(8):e0200061. doi: 10.1371/journal.pone.0200061.

36: Sutcliffe BK, Bennett PN, Fraser SF, Mohebbi M. The deterioration in physical function of hemodialysis patients. *Hemodial Int*. 2018 Apr;22(2):245-253. doi: 10.1111/hdi.12570.

37: Suzuki T, Ikeda M, Minami M, Matayoshi Y, Nakao M, Nakamura T, Abo M. Beneficial Effect of Intradialytic Electrical Muscle Stimulation in Hemodialysis Patients: A Randomized Controlled Trial. *Artif Organs*. 2018 Sep;42(9):899-910. doi: 10.1111/aor.13161.

38: Torino C, Panuccio V, Tripepi R, Vilasi A, Postorino M, Tripepi G, Mallamaci F, Zoccali C. The dominant prognostic value of physical functioning among quality of life domains in end-stage kidney disease. *Nephrol Dial Transplant*. 2018 Nov 15. doi: 10.1093/ndt/gfy334.

39: Valenzuela PL, de Alba A, Pedrero-Chamizo R, Morales JS, Cobo F, Botella A, González-Gross M, Pérez M, Lucia A, Marín-López MT. Intradialytic Exercise: One Size Doesn't Fit All. *Front Physiol*. 2018 Jul 5;9:844. doi:10.3389/fphys.2018.00844.

40: Wilschut ED, Rotmans JI, Bos EJ, van Zoest D, Eefting D, Hamming JF, van der Bogt KEA. Supervised preoperative forearm exercise to increase blood vessel diameter in patients requiring an arteriovenous access for hemodialysis: rationale and design of the PINCH trial. *J Vasc Access*. 2018 Jan;19(1):84-88. doi: 10.5301/jva.5000826.

41: Yamamoto S, Matsuzawa R, Abe Y, Hoshi K, Yoneki K, Harada M, Watanabe T, Shimoda T, Suzuki Y, Matsunaga Y, Kamiya K, Yoshida A, Matsunaga A. Utility of Regular Management of Physical Activity and Physical Function in Hemodialysis Patients. *Kidney Blood Press Res.* 2018;43(5):1505-1515. doi: 10.1159/000494016.

42: Young HML, Jeurkar S, Churchward DR, Dungey M, Stensel DJ, Bishop NC, Greenwood SA, Singh SJ, Smith AC, Burton JO. Implementing a theory-based intradialytic exercise programme in practice: a quality improvement project. *Clin Kidney J.* 2018 Dec;11(6):832-840. doi: 10.1093/ckj/sfy050.

43: Young HML, March DS, Graham-Brown MPM, Jones AW, Curtis F, Grantham CS, Churchward DR, Highton P, Smith AC, Singh SJ, Bridle C, Burton JO. Effects of intradialytic cycling exercise on exercise capacity, quality of life, physical function and cardiovascular measures in adult haemodialysis patients: a systematic review and meta-analysis. *Nephrol Dial Transplant.* 2018 Aug 1;33(8):1436-1445. doi: 10.1093/ndt/gfy045.

44: Ünver S, Akyolcu N. The Effect of Hand Exercise on Reducing the Symptoms in Hemodialysis Patients with Carpal Tunnel Syndrome. *Asian J Neurosurg.* 2018 Jan-Mar;13(1):31-36. doi: 10.4103/ajns.AJNS_343_16.