

Bullani R, El-Housseini Y, Giordano F, Larcinese A, Ciutto L, Bertrand PC, Wuerzner G, Burnier M, Teta D. Effect of intradialytic resistance band exercise on physical function in patients on maintenance hemodialysis: a pilot study. <i>J Ren Nutr.</i> 2011 Jan;21(1):61-5. PubMed PMID: 21195922.
Cheema, B. S., H. Abas, et al.: Effect of resistance training during hemodialysis on circulating cytokines: a randomized controlled trial. <i>Eur J Appl Physiol</i> 2011; 7: 1437-1445.
Clapp EL, Bevington A, Smith AC. Exercise for children with chronic kidneydisease and end-stage renal disease. <i>Pediatr Nephrol.</i> 2011 Jan 14. [Epub ahead of print] PubMed PMID: 21229267.
De Lima MC, Cicotoste Cde I, Cardoso Kda S, et al: Effect of exercise performed during hemodialysis: strength versus aerobic. <i>Ren Fail</i> 2013; 35: 697-704
Delgado C1, Johansen KL: Barriers to exercise participation among dialysis patients. <i>Nephrol Dial Transplant.</i> 2012; 3:1152-7. doi: 10.1093/ndt/gfr404. Epub 2011 Jul 26.
Fotbolcu H, Duman D, Ecder SA, et al: Attenuated cardiovascular response to sympathetic system activation during exercise in patients with dialysis induced hypotension. <i>Am J Nephrol</i> 2011; 33: 491-498
Garibotto G, Sofia A, Saffioti S, Bonanni A, Mannucci I, Parodi EL, Cademartori V, Verzola D. Effects of peritoneal dialysis on protein metabolism. <i>Nutr Metab Cardiovasc Dis.</i> 2012 Aug 13. [Epub ahead of print] PubMed PMID: 22898450.
Giannaki, C. D., I. Stefanidis, et al. (2011). "The effect of prolonged intradialytic exercise in hemodialysis efficiency indices." <i>ASAIO J</i> 2011; 3: 213-218.
Giannaki, CD, Hadjigeorgiou GM, Karatzafiri C, et al: A single-blind randomized controlled trial to evaluate the effect of 6 month of progressive aerobic exercise training in patients with uraemic restless legs syndrome. <i>Nephrol Dial Transplant</i> 2013; 28: 2834-2840
Giannaki, CD, Sakkas GK, Karatzafiri C, et al: Effect of exercise training and dopamine agonists in patients with uremicrestless legs syndrome: a six-month randomized, partially double-blind, placebo-controlled comparative study. <i>BMC Nephrol</i> 2013; 14: 194
Gregory SM, Headley SA, Germain M, Flyvbjerg A, Frystyk J, Coughlin MA, Milch CM, Sullivan S, Nindl BC. Lack of circulating bioactive and immunoreactive IGF-I changes despite improved fitness in chronic kidney disease patients following 48 weeks of physical training. <i>Growth Horm IGF Res.</i> 2011 Feb;21(1):51-6. Epub 2011Jan 19. PubMed PMID: 21251861.
Heiwe S, Jacobson SH: Exercise training for adults with chronic kidney disease. <i>Cochrane Database Syst Rev</i> 2011; 10:CD003236
Ikizler TA. Exercise as an anabolic intervention in patients with end-stage renal disease. <i>J Ren Nutr.</i> 2011 Jan;21(1):52-6. Review. PubMed PMID: 21195920; PubMed Central PMCID: PMC3061820.
Koufaki P, Greenwood SA, Macdougall IC, Mercer TH: Exercise therapy in individuals with chronic kidney disease: a systematic review and synthesis of the research evidence. <i>Annu Rev Nurs Res.</i> 2013;31:235-75. doi: 10.1891/0739-6686.31.235.
Makhlough A, Ilali E, Mohseni R, Shamohammadi S: Effect of intradialytic aerobic exercise on serum elcktroytes levels in hemodialysis patients.Iran J Kidney Dis 2013;6:119-123
Mohseni R, Emami Zeydi A, Illali E, Adib-Hajbaghery M, Makhlough A: The effect of intradialytic aerobic exercise on dialysis efficacy in hemodialysis patients: a randomized controlled trial. <i>Oman Med J</i> 2013;28:345-349
Oishi D, Koitabashi K, Hiraki K, Imai N, Sakurada T, Konno Y, Shibagaki Y, Yasuda T, Kimura K. Physical activity is associated with serum albumin in peritoneal dialysis patients. <i>Adv Perit Dial.</i> 2012;28:148-52. PubMed PMID: 23311233.
Painter P, Krasnoff JB, Kuskowski M, Frassetto L, Johansen KL. Effects of modality change and transplant on peak oxygen uptake in patients with kidney failure. <i>Am J Kidney Dis.</i> 2011 Jan;57(1):113-22. Epub 2010 Sep 25. PubMed PMID: 20870330; PubMed Central PMCID: PMC3010466.
Pellizzaro CO, Thome FS, Veronese FV: Effect of peripheral and respiratory muscle training on the functional capacity of hemodialysis patients. <i>Ren Fail</i> 2013;35:189-197
Schaar, B., M. Feldkotter, et al. (2011). "Cardiorespiratory capacity in children and adolescents on maintenance haemodialysis." <i>Nephrol Dial Transplant</i> 26(11): 3701-3708.
SmartN, Steele M: Exercise training in haemodialysis patients: a stematic review and metaanalysis. <i>Nephrology (Carlton)</i> 2011;16:626-632
Song WJ, Sohng KY: Effects of progressive resistance training on body composition, physical fitness and quality of life of patients on hemodialysis. <i>J Korean Acad Nurs</i> 2012;42:947-956